

# Kansas Agricultural Experiment Station Research Reports

Volume 1  
Issue 6 *Turfgrass Research*

Article 13

January 2015

## Crabgrass Control in Spring Seeding Using Early and Late Combinations of Prodiamine and Dithiopyr

J. Hoyle

Kansas State University, [jahoyle@ksu.edu](mailto:jahoyle@ksu.edu)

J. Reeves

Kansas State University, [jakereeves@ksu.edu](mailto:jakereeves@ksu.edu)

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Horticulture Commons](#), and the [Weed Science Commons](#)

### Recommended Citation

Hoyle, J. and Reeves, J. (2015) "Crabgrass Control in Spring Seeding Using Early and Late Combinations of Prodiamine and Dithiopyr," *Kansas Agricultural Experiment Station Research Reports*: Vol. 1: Iss. 6. <https://doi.org/10.4148/2378-5977.1101>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright January 2015 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



---

## Crabgrass Control in Spring Seeding Using Early and Late Combinations of Prodiamine and Dithiopyr

### Abstract

With less crabgrass pressure, the Kansas site showed an inhibitory effect of pre-emergent herbicides on spring tall fescue (*Lolium arundinaceum*) seeding and establishment. However, at the Nebraska site, greater crabgrass pressure showed a benefit of pre-emergent herbicide use in new spring seeding of tall fescue.

### Keywords

turfgrass, *Digitaria* spp., crabgrass control, prodiamine, dithiopyr, pre-emergent herbicide, spring tall fescue, *Lolium arundinacea*

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

# TURFGRASS RESEARCH 2015



JULY 2015



Kansas State University  
Agricultural Experiment Station  
and Cooperative Extension Service

K-State Research and Extension is an equal  
opportunity provider and employer.

## Crabgrass Control in Spring Seeding Using Early and Late Combinations of Prodiamine and Dithiopyr

*Jared Hoyle and Jake Reeves*

*Collaborators: Zac Reicher and Matthew Sousek, University of Nebraska-Lincoln*

**Summary.** With less crabgrass pressure, the Kansas site showed an inhibitory effect of pre-emergent herbicides on spring tall fescue (*Lolium arundinaceum*) seeding and establishment. However, at the Nebraska site, greater crabgrass pressure showed a benefit of pre-emergent herbicide use in new spring seeding of tall fescue.

**Rationale.** Crabgrass (*Digitaria* spp.) germination typically occurs during cool-season turfgrass spring seeding. Pre-emergent herbicides are recommended for control of summer annual grassy weeds such as crabgrass but can inhibit germination of desired turfgrass. Reduced rates with split applications of dithiopyr and prodiamine in liquid and dry formulations could decrease crabgrass pressure without inhibiting turfgrass establishment.

**Objective.** Determine if various application timings or rates would improve seedling safety or crabgrass control.

**Study Description.** Research was conducted at the Rocky Ford Turfgrass Research Center in Manhattan, Kansas, and the John Seaton Anderson Turfgrass Research Center in Mead, Nebraska. Research sites were treated with Roundup+Reward 2 weeks before seeding. Plots were established by seeding tall fescue at 10 lb/1,000 ft<sup>2</sup>, followed by the initial application of herbicides, power-raking in two directions, and an application of 11-52-0 at 1 lb P<sub>2</sub>O<sub>5</sub>/1,000 ft<sup>2</sup>. This experiment was designed as a 2 x 2 x 6 factorial with two active ingredients, two formulations each, and six application strategies. Plots were 5 x 5 ft in each location with three replications.

View all turfgrass research reports online at: <http://newprairiepress.org/kaesrr>





Applications were made with a CO<sub>2</sub> backpack sprayer using a three-nozzle boom with 8002VS nozzles at 30 psi in 2 gal/1,000 ft<sup>2</sup>, or spread with a shaker can. Areas were mowed at 2.5 inches once seedlings reached approximately 3 inches and weekly thereafter. Cover ratings of tall fescue, large crabgrass, and yellow foxtail were taken at 2-week intervals. This experiment was designed as a 2 x 2 x 6 factorial with two active ingredients, two formulations each, and six application strategies. Means were separated using Fisher's Protected LSD ( $P \leq 0.05$ ).

**Results.** Dithiopyr appeared to be slightly safer than prodiamine for this use, and applications 4 weeks after seeding appeared to be safer on tall fescue and as effective for crabgrass control as at-seeding applications. Granular fertilizer formulations appeared to be safer on tall fescue than sprayable formulations. Outside stresses such as drought or high temperatures may exacerbate any possible damage.



**Figure 1.** Research trial location at Rocky Ford Turfgrass Research Center in Manhattan, Kansas.



Kansas State University  
Agricultural Experiment Station  
and Cooperative Extension Service

